

Title	N- Fractional Calculus and n th Derivatives of Some Logarithmic Functions (Study on Non-Analytic and Univalent Functions and Applications)
Author(s)	Romero, Susana S. de; Nishimoto, Katsuyuki
Citation	数理解析研究所講究録 (2009), 1626: 99-108
Issue Date	2009-01
URL	http://hdl.handle.net/2433/140312
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

N- Fractional Calculus and $n(\in \mathbb{Z}^+)$ th Derivatives of Some Logarithmic Functions

* Susana S. de Romero and ** Katsuyuki Nishimoto

* CIMA, Facultad de Ingenieria, Universidad del Zulia,
Apartado 10482, Maracaibo, VENEZUELA.

** Institute for Applied Mathematics, Descartes Press Co.
2 - 13 - 10 Kaguike, Koriyama 963 - 8833, JAPAN.

Abstract

In this article N- fractional calculus and n -th derivatives of logarithmic functions

$$\log((z-b)^m - c), \quad (m \in \mathbb{Z}_0^+) \quad \text{and} \quad \log(z^2 + 2az + d)$$

are reported.

That is, we have the following, for example.

$$(i) \quad (\log((z-b)^m - c))_\gamma = -e^{-i\pi\gamma} m(z-b)^{-\gamma} \Gamma(\gamma)$$

$$\times \sum_{k=0}^{\infty} \frac{\Gamma(mk + \gamma)}{\Gamma(\gamma)\Gamma(mk + 1)} \left(\frac{c}{(z-b)^m} \right)^k$$

$$(|\Gamma(\gamma)|, |\Gamma(mk + \gamma)| < \infty),$$

and

$$(ii) \quad (\log((z-b)^m - c))_n = (-1)^{n+1} m(z-b)^{-n} \Gamma(n)$$

$$\times \sum_{k=0}^{\infty} \frac{\Gamma(mk + n)}{\Gamma(n)\Gamma(mk + 1)} \left(\frac{c}{(z-b)^m} \right)^k$$

$$(n \in \mathbb{Z}^+),$$

where

$$(z-b)^m - c \neq 0, 1, \quad \text{and} \quad |c/(z-b)^m| < 1.$$

§ 0. Introduction (Definition of Fractional Calculus)

(I) Definition. (by K. Nishimoto) ([1] Vol. 1)

Let $D = \{D_-, D_+\}$, $C = \{C_-, C_+\}$,

C_- be a curve along the cut joining two points z and $-\infty + i\text{Im}(z)$,

C_+ be a curve along the cut joining two points z and $\infty + i\text{Im}(z)$,

D_- be a domain surrounded by C_- , D_+ be a domain surrounded by C_+ .

(Here D contains the points over the curve C).

Moreover, let $f = f(z)$ be a regular function in $D(z \in D)$,

$$f_v = (f)_v = {}_C(f)_v = \frac{\Gamma(v+1)}{2\pi i} \int_C \frac{f(\xi)}{(\xi-z)^{v+1}} d\xi \quad (v \notin \mathbb{Z}), \quad (1)$$

$$(f)_{-m} = \lim_{v \rightarrow -m} (f)_v \quad (m \in \mathbb{Z}^+), \quad (2)$$

where $-\pi \leq \arg(\xi-z) \leq \pi$ for C_- , $0 \leq \arg(\xi-z) \leq 2\pi$ for C_+ ,

$\xi \neq z$, $z \in \mathbb{C}$, $v \in \mathbb{R}$, Γ ; Gamma function,

then $(f)_v$ is the fractional differintegration of arbitrary order v (derivatives of order v for $v > 0$, and integrals of order $-v$ for $v < 0$), with respect to z , of the function f , if $|(f)_v| < \infty$.

(II) On the fractional calculus operator N^v [3]

Theorem A. Let fractional calculus operator (Nishimoto's Operator) N^v be

$$N^v = \left(\frac{\Gamma(v+1)}{2\pi i} \int_C \frac{d\xi}{(\xi-z)^{v+1}} \right) \quad (v \notin \mathbb{Z}), \quad [\text{Refer to (1)}] \quad (3)$$

with
$$N^{-m} = \lim_{v \rightarrow -m} N^v \quad (m \in \mathbb{Z}^+), \quad (4)$$

and define the binary operation \circ as

$$N^\beta \circ N^\alpha f = N^\beta N^\alpha f = N^\beta (N^\alpha f) \quad (\alpha, \beta \in \mathbb{R}), \quad (5)$$

then the set

$$\{N^v\} = \{N^v | v \in \mathbb{R}\} \quad (6)$$

is an Abelian product group (having continuous index v) which has the inverse transform operator $(N^v)^{-1} = N^{-v}$ to the fractional calculus operator N^v , for the

N- Fractional Calculus and $n (\in \mathbb{Z}^+)$ th Derivatives of Some Logarithmic Functions

function f such that $f \in F = \{f; 0 \neq |f_v| < \infty, v \in \mathbb{R}\}$, where $f = f(z)$ and $z \in \mathbb{C}$.
(vis. $-\infty < v < \infty$).

(For our convenience, we call $N^\beta \circ N^\alpha$ as product of N^β and N^α .)

Theorem B. "F.O.G. $\{N^\nu\}$ " is an "Action product group which has continuous index ν " for the set of F . (F.O.G.; Fractional calculus operator group) [3]

Theorem C. Let

$$S := \{\pm N^\nu\} \cup \{0\} = \{N^\nu\} \cup \{-N^\nu\} \cup \{0\} \quad (\nu \in \mathbb{R}). \quad (7)$$

Then the set S is a commutative ring for the function $f \in F$, when the identity

$$N^\alpha + N^\beta = N^\gamma \quad (N^\alpha, N^\beta, N^\gamma \in S) \quad (8)$$

holds. [5]

(III) **Lemma.** We have [1]

$$(i) \quad ((z-c)^b)_\alpha = e^{-i\pi\alpha} \frac{\Gamma(\alpha-b)}{\Gamma(-b)} (z-c)^{b-\alpha} \quad \left(\left| \frac{\Gamma(\alpha-b)}{\Gamma(-b)} \right| < \infty \right),$$

$$(ii) \quad (\log(z-c))_\alpha = -e^{-i\pi\alpha} \Gamma(\alpha) (z-c)^{-\alpha} \quad (|\Gamma(\alpha)| < \infty),$$

$$(iii) \quad ((z-c)^{-\alpha})_{-\alpha} = -e^{i\pi\alpha} \frac{1}{\Gamma(\alpha)} \log(z-c) \quad (|\Gamma(\alpha)| < \infty),$$

where $z-c \neq 0$ for (i) and $z-c \neq 0, 1$ for (ii), (iii),

§ 1. Preliminary

Theorem D. below for the fractional calculus of a logarithmic function is reported by K. Nishimoto (cf. J. Frac. Calc. Vol. 29, May (2006), p. 40.).

Theorem D. We have

$$(i) \quad (\log((z-b)^\beta - c))_\gamma = -e^{-i\pi\gamma} \beta (z-b)^{-\gamma} \Gamma(\gamma)$$

$$\times \sum_{k=0}^{\infty} \frac{\Gamma(\beta k + \gamma)}{\Gamma(\gamma) \Gamma(\beta k + 1)} \left(\frac{c}{(z-b)^\beta} \right)^k \quad (|\Gamma(\gamma)|, \left| \frac{\Gamma(\beta k + \gamma)}{\Gamma(\beta k + 1)} \right| < \infty), \quad (1)$$

and

$$(ii) \quad (\log((z-b)^\beta - c))_m = (-1)^{m+1} \beta (z-b)^{-m} \Gamma(m)$$

$$\times \sum_{k=0}^{\infty} \frac{\Gamma(\beta k + m)}{\Gamma(m) \Gamma(\beta k + 1)} \left(\frac{c}{(z-b)^\beta} \right)^k \quad (m \in \mathbb{Z}^+), \quad (2)$$

where

$$(z-b)^\beta - c \neq 0, 1, \quad \text{and} \quad |c/(z-b)^\beta| < 1.$$

Susana S. de Romero and Katsuyuki Nishimoto

§ 2. N-Fractional Calculus of Functions

$$\log((z-b)^m - c)$$

Theorem 1. *We have*

$$\begin{aligned} (i) \quad & (\log((z-b)^m - c))_\gamma = -e^{-i\pi\gamma} m(z-b)^{-\gamma} \Gamma(\gamma) \\ & \times \sum_{k=0}^{\infty} \frac{\Gamma(mk + \gamma)}{\Gamma(\gamma)\Gamma(mk + 1)} \left(\frac{c}{(z-b)^m} \right)^k \quad (1) \\ & (|\Gamma(\gamma)|, |\Gamma(mk + \gamma)| < \infty), \end{aligned}$$

and

$$\begin{aligned} (ii) \quad & (\log((z-b)^m - c))_n = (-1)^{n+1} m(z-b)^{-n} \Gamma(n) \\ & \times \sum_{k=0}^{\infty} \frac{\Gamma(mk + n)}{\Gamma(n)\Gamma(mk + 1)} \left(\frac{c}{(z-b)^m} \right)^k \quad (n \in \mathbb{Z}^+), \quad (2) \end{aligned}$$

where

$$m \in \mathbb{Z}_0^+, \quad (z-b)^m - c \neq 0, 1, \quad \text{and} \quad |c/(z-b)^m| < 1.$$

Proof of (i). Set $\beta = m$ in Theorem D. (i) in Preliminary we have (1) clearly, under the conditions stated before.

Proof of (ii). Set $\gamma = n$ in (1), we have then (2).

Corollary 1. *We have*

$$\begin{aligned} (i) \quad & (\log((z-b)^2 - c))_\gamma = -e^{-i\pi\gamma} 2(z-b)^{-\gamma} \Gamma(\gamma) \\ & \times \sum_{k=0}^{\infty} \frac{\Gamma(2k + \gamma)}{\Gamma(\gamma)\Gamma(2k + 1)} \left(\frac{c}{(z-b)^2} \right)^k \quad (3) \\ & (|\Gamma(\gamma)|, |\Gamma(2k + \gamma)| < \infty), \end{aligned}$$

and

$$\begin{aligned} (ii) \quad & (\log((z-b)^2 - c))_n = (-1)^{n+1} 2(z-b)^{-n} \Gamma(n) \\ & \times \sum_{k=0}^{\infty} \frac{\Gamma(2k + n)}{\Gamma(n)\Gamma(2k + 1)} \left(\frac{c}{(z-b)^2} \right)^k \quad (n \in \mathbb{Z}^+), \quad (4) \end{aligned}$$

where

$$(z-b)^2 - c \neq 0, 1, \quad \text{and} \quad |c/(z-b)^2| < 1.$$

Proof. Set $m = 2$ in Theorem 1.

N- Fractional Calculus and $n (\in \mathbb{Z}^+)$ th Derivatives of Some Logarithmic Functions

Theorem 2. We have the following identity.

$$(i) \quad 2 \sum_{k=0}^{\infty} \frac{\Gamma(2mk + \gamma)}{\Gamma(\gamma)\Gamma(2mk + 1)} T^k = \sum_{k=0}^{\infty} \frac{\Gamma(mk + \gamma)}{\Gamma(\gamma)\Gamma(mk + 1)} \{(T^{1/2})^k + (-T^{1/2})^k\} \quad (5)$$

$$(|\Gamma(mk + \gamma)| < \infty),$$

and

$$(ii) \quad 2 \sum_{k=0}^{\infty} \frac{[2mk + 1]_{n-1}}{\Gamma(n)} T^k = \sum_{k=0}^{\infty} \frac{[mk + 1]_{n-1}}{\Gamma(n)} \{(T^{1/2})^k + (-T^{1/2})^k\} \quad (6)$$

$$(n \in \mathbb{Z}^+),$$

where

$$T = \frac{c}{(z-b)^{2m}}, \quad |T| < 1, \quad m \in \mathbb{Z}^+, \quad (7)$$

$$[\lambda]_k = \lambda(\lambda + 1) \cdots (\lambda + k - 1) = \Gamma(\lambda + k) / \Gamma(\lambda) \text{ with } [\lambda]_0 = 1.$$

(Notation of Pochhammer).

Proof of (i). We have

$$\log((z-b)^{2m} - c) = \log((z-b)^m - \sqrt{c}) + \log((z-b)^m + \sqrt{c}). \quad (8)$$

$$((z-b)^m \pm \sqrt{c} \neq 0, 1).$$

Operate N^γ to the both sides of (8), we have then

$$(\log((z-b)^{2m} - c))_\gamma = (\log((z-b)^m - \sqrt{c}))_\gamma + (\log((z-b)^m + \sqrt{c}))_\gamma. \quad (9)$$

Now we have

$$(\log((z-b)^{2m} - c))_\gamma = -e^{-i\pi\gamma} 2m(z-b)^{-\gamma} \Gamma(\gamma) \sum_{k=0}^{\infty} \frac{\Gamma(2mk + \gamma)}{\Gamma(\gamma)\Gamma(2mk + 1)} T^k \quad (10)$$

$$(|\Gamma(2mk + \gamma)| < \infty),$$

and

$$(\log((z-b)^m - \sqrt{c}))_\gamma = -e^{-i\pi\gamma} m(z-b)^{-\gamma} \Gamma(\gamma) \sum_{k=0}^{\infty} \frac{\Gamma(mk + \gamma)}{\Gamma(\gamma)\Gamma(mk + 1)} (T^{1/2})^k \quad (11)$$

$$(|\Gamma(mk + \gamma)| < \infty),$$

by Theorem D. (i), respectively.

Susana S. de Romero and Katsuyuki Nishimoto

Then we obtain (i) from (9), applying (10) and (11), under the conditions.

Proof of (ii). Set $\gamma = n$ in (i).

Corollary 2. We have the following identity.

$$(i) \quad 2(z-b)^{-\gamma} \sum_{k=0}^{\infty} \frac{\Gamma(2k+\gamma)}{\Gamma(\gamma)\Gamma(2k+1)} T^k = \frac{1}{(z-b-\sqrt{c})^\gamma} + \frac{1}{(z-b+\sqrt{c})^\gamma} \quad (12)$$

$$(\quad |\Gamma(2k+\gamma)| < \infty),$$

and

$$(ii) \quad 2(z-b)^{-n} \sum_{k=0}^{\infty} \frac{[2k+1]_{n-1}}{\Gamma(n)} T^k = \frac{1}{(z-b-\sqrt{c})^n} + \frac{1}{(z-b+\sqrt{c})^n} \quad (13)$$

$$(n \in \mathbb{Z}^+).$$

Proof of (i). Set $m = 1$ in Theorem 2. (i), we have then

$$2 \sum_{k=0}^{\infty} \frac{\Gamma(2k+\gamma)}{\Gamma(\gamma)\Gamma(2k+1)} T^k = \sum_{k=0}^{\infty} \frac{\Gamma(k+\gamma)}{\Gamma(\gamma)\Gamma(k+1)} \{(T^{1/2})^k + (-T^{1/2})^k\}. \quad (14)$$

Now we have

$$\sum_{k=0}^{\infty} \frac{\Gamma(k+\gamma)}{\Gamma(\gamma)\Gamma(k+1)} (T^{1/2})^k = \sum_{k=0}^{\infty} \frac{[\gamma]_k}{k!} (T^{1/2})^k \quad (15)$$

$$= \frac{(z-b)^\gamma}{(z-b-\sqrt{c})^\gamma}. \quad (16)$$

Therefore, we obtain (12) from (14) and (16).

Proof of (ii). Set $\gamma = n$ in (i).

Note. Other proof of (i). We have

$$(\log((z-b)^2 - c))_\gamma = (\log((z-b) - \sqrt{c}))_\gamma + (\log((z-b) + \sqrt{c}))_\gamma, \quad (17)$$

from (9), setting $m = 1$.

Next we have

$$(\log((z-b) - \sqrt{c}))_\gamma = -e^{-i\pi\gamma} \Gamma(\gamma) (z-b-\sqrt{c})^{-\gamma}, \quad (|\Gamma(\gamma)| < \infty) \quad (18)$$

by Lemma (ii).

Therefore, we obtain (12) from (17), applying (3) and (18).

N- Fractional Calculus and $n(\in \mathbb{Z}^+)$ th Derivatives of Some Logarithmic Functions

Theorem 3. We have

$$(i) \quad (\log(z^2 + 2az + d))_\gamma = -e^{-i\pi\gamma} 2(z+a)^{-\gamma} \Gamma(\gamma) \\ \times \sum_{k=0}^{\infty} \frac{\Gamma(2k+\gamma)}{\Gamma(\gamma)\Gamma(2k+1)} \left(\frac{a^2-d}{(z+a)^2} \right)^k \quad (19) \\ (|\Gamma(\gamma)|, |\Gamma(2k+\gamma)| < \infty),$$

and

$$(ii) \quad (\log(z^2 + 2az + d))_n = (-1)^{n+1} 2(z+a)^{-n} \Gamma(n) \\ \times \sum_{k=0}^{\infty} \frac{\Gamma(2k+n)}{\Gamma(n)\Gamma(2k+1)} \left(\frac{a^2-d}{(z+a)^2} \right)^k \quad (n \in \mathbb{Z}^+), \quad (20)$$

where

$$z^2 + 2az + d \neq 0, 1, \quad \text{and} \quad |(a^2-d)/(z+a)^2| < 1.$$

Proof of (i). We have

$$z^2 + 2az + d = (z+a)^2 - c, \quad (c = a^2 - d). \quad (21)$$

hence

$$\log(z^2 + 2az + d) = \log((z+a)^2 - c). \quad (22)$$

Operate N^γ to the both sides of (22), we have then

$$(\log(z^2 + 2az + d))_\gamma = (\log((z+a)^2 - c))_\gamma, \quad (23)$$

therefore, we obtain (19) clearly, setting $b = -a$ and $c = a^2 - d$ in Corollary 1.

(i), under the conditions stated before.

Proof of (ii). Set $\gamma = n$ in (19).

§ 3. Semi Derivatives and Integrals

Corollary 3. We have

$$(i) \quad (\log((z-b)^m - c))_{1/2} = im(z-b)^{-1/2} \sqrt{\pi} \\ \times \sum_{k=0}^{\infty} \frac{\Gamma(mk + \frac{1}{2})}{\Gamma(\frac{1}{2})\Gamma(mk+1)} \left(\frac{c}{(z-b)^m} \right)^k \quad (1) \\ (\text{semi derivatives}),$$

and

Susana S. de Romero and Katsuyuki Nishimoto

$$\begin{aligned}
 (ii) \quad & (\log((z-b)^m - c))_{-1/2} = im(z-b)^{1/2} 2\sqrt{\pi} \\
 & \times \sum_{k=0}^{\infty} \frac{\Gamma(mk - \frac{1}{2})}{\Gamma(-\frac{1}{2})\Gamma(mk+1)} \left(\frac{c}{(z-b)^m} \right)^k \quad (2) \\
 & \text{(semi integrals),}
 \end{aligned}$$

where

$$m \in \mathbb{Z}_0^+, \quad (z-b)^m - c \neq 0, 1, \quad \text{and} \quad |c/(z-b)^m| < 1.$$

Proof. Set $\gamma = 1/2$ and $-1/2$ in Theorem 1.(i), we have then (1) and (2) respectively

Corollary 4. We have

$$\begin{aligned}
 (i) \quad & (\log((z-b)^2 - c))_{1/2} = i2\sqrt{\pi}(z-b)^{-1/2} \sum_{k=0}^{\infty} \frac{\Gamma(2k + \frac{1}{2})}{\Gamma(\frac{1}{2})\Gamma(2k+1)} \left(\frac{c}{(z-b)^2} \right)^k \quad (3) \\
 & \text{(semi derivatives),}
 \end{aligned}$$

and

$$\begin{aligned}
 (ii) \quad & (\log((z-b)^2 - c))_{-1/2} = i4\sqrt{\pi}(z-b)^{1/2} \sum_{k=0}^{\infty} \frac{\Gamma(2k - \frac{1}{2})}{\Gamma(-\frac{1}{2})\Gamma(2k+1)} \left(\frac{c}{(z-b)^2} \right)^k \quad (4) \\
 & \text{(semi integrals),}
 \end{aligned}$$

where

$$(z-b)^2 - c \neq 0, 1, \quad \text{and} \quad |c/(z-b)^2| < 1.$$

Proof. Set $m = 2$ in Corollary 3.

Corollary 5. We have

$$\begin{aligned}
 (i) \quad & (\log((z-b)^2 - c))_{1/2} = i\sqrt{\pi} \left(\frac{1}{\sqrt{z-b-\sqrt{c}}} + \frac{1}{\sqrt{z-b+\sqrt{c}}} \right) \quad (5) \\
 & \text{(semi derivatives)}
 \end{aligned}$$

and

$$\begin{aligned}
 (i) \quad & (\log((z-b)^2 - c))_{-1/2} = i2\sqrt{\pi} \left(\sqrt{z-b-\sqrt{c}} + \sqrt{z-b+\sqrt{c}} \right) \quad (6) \\
 & \text{(semi integrals)}
 \end{aligned}$$

where

$$(z-b)^2 - c \neq 0, 1.$$

Proof. Set $\gamma = 1/2$ and $-1/2$ in Corollary 2, we have then (5) and (6) respectively, under the conditions.

§ 4. Some Special Cases

[I] When $n = 1$, we have

$$(\log((z-b)^2 - c))_1 = 2(z-b)^{-1} \sum_{k=0}^{\infty} \left(\frac{c}{(z-b)^2} \right)^k \quad (1)$$

$$= 2(z-b)^{-1} \sum_{k=0}^{\infty} \frac{[1]_k}{k!} \left(\frac{c}{(z-b)^2} \right)^k = 2(z-b)^{-1} \left(1 - \frac{c}{(z-b)^2} \right)^{-1} \quad (2)$$

$$= 2((z-b)^2 - c)^{-1}(z-b) \quad , \quad (3)$$

from Corollary 1 (ii).

[II] When $n = 2$, we have

$$(\log((z-b)^2 - c))_2 = -2(z-b)^{-2} \sum_{k=0}^{\infty} \frac{\Gamma(2k+2)}{\Gamma(2k+1)} \left(\frac{c}{(z-b)^2} \right)^k \quad (4)$$

$$= -2(z-b)^{-2} \sum_{k=0}^{\infty} \frac{[1]_k (2k+1)}{k!} T^k \quad \left(T = \frac{c}{(z-b)^2} \right) \quad (5)$$

$$= -2(z-b)^{-2} \left\{ \sum_{k=0}^{\infty} \frac{[1]_k 2k}{k!} T^k + \sum_{k=0}^{\infty} \frac{[1]_k}{k!} T^k \right\} \quad (6)$$

$$= 2(z-b)^{-2} \{ 2T(1-T)^{-2} + (1-T)^{-1} \} \quad (7)$$

$$= -2((z-b)^2 + c) ((z-b)^2 - c)^{-2} \quad (8)$$

from Corollary 1 (ii).

This result coincides with the one obtained by the classical calculus ;

$$\frac{d^2}{dz^2} \log((z-b)^2 - c) \quad .$$

Note.

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{[1]_k 2k}{k!} T^k &= 2 \sum_{k=1}^{\infty} \frac{[1]_k}{(k-1)!} T^k = 2T \sum_{k=0}^{\infty} \frac{[1]_{k+1}}{k!} T^k \\ &= 2T \sum_{k=0}^{\infty} \frac{[2]_k}{k!} T^k = 2T(1-T)^{-2} . \end{aligned}$$

Susana S. de Romero and Katsuyuki Nishimoto

References

- [1] K. Nishimoto ; Fractional Calculus, Vol. 1 (1984), Vol. 2 (1987), Vol. 3 (1989), Vol. 4 (1991), Vol. 5, (1996), Descartes Press, Koriyama, Japan.
- [2] K. Nishimoto ; An Essence of Nishimoto's Fractional Calculus (Calculus of the 21st Century); Integrals and Differentiations of Arbitrary Order (1991), Descartes Press, Koriyama, Japan.
- [3] K. Nishimoto ; On Nishimoto's fractional calculus operator N^ν (On an action group), J. Frac. Calc. Vol. 4, Nov. (1993), 1 - 11.
- [4] K. Nishimoto ; Unification of the integrals and derivatives (A serendipity in fractional calculus), J. Frac. Calc. Vol. 6, Nov. (1994), 1 - 14.
- [5] K. Nishimoto ; Ring and Field Produced from The Set of N- Fractional Calculus Operator, J. Frac Calc. Vol. 24, Nov. (2003), 29 - 36.
- [6] K. Nishimoto ; On the fractional calculus $(a - z)^\beta$ and $\log(a - z)$, J. Frac. Calc. Vol.3, May (1993), 19 - 27.
- [7] K. Nishimoto and S.- T. Tu ; Fractional calculus of Psi functions (Generalized Polygamma unctions), J. Frac. Calc. Vol.5 May (1994), 27 - 34.
- [8] S.- T. Tu and K. Nishimoto ; On the fractional calculus of functions $(cz - a)^\beta \log(cz - a)$, J. Frac.Calc.Vol.5, May (1994), 35 - 43.
- [9] K. Nishimoto ; N- Fractional Calculus of the Power and Logarithmic Functions and Some Identities, J. Frac. Calc. Vol.21, May (2002), 1 - 6.
- [10] K. Nishimoto ; Some Theorems for N- Fractional Calculus of Logarithmic Functions I, J. Frac Calc.Vol.21, May (2002), 7 - 12.
- [11] K. Nishimoto ; N- Fractional Calculus of Products of Some Power Functions, J. Frac.Calc. Vol.27, May (2005), 83 - 88.
- [12] K. Nishimoto ; N- Fractional Calculus of Some Composite Functions, J. Frac. Calc. Vol. 29, May (2006), 35 - 44.
- [13] K. Nishimoto ; N- Fractional Calculus of Some Composite Algebraic Functions, J. Frac.Calc. Vol. 31, May (2006), 11 - 23.
- [14] K. Nishimoto ; N- Fractional Calculus of Some Elementary Functions and Their Semi Differintegrations, J. Frac. Caic. Vol. 31, May (2007), 1 - 10.
- [15] K. Nishimoto and T. Miyakoda ; N- Fractional Calculus and n -th Derivatives of Some Algebraic Functions, J. Frac. Calc. Vol. 31, May (2007), 53 - 62.
- [16] T. Miyakoda ; N- Fractional Calculus of Certain Algebraic Functions, J. Frac. Calc.Vol. 31, May (2007), 63 - 76.
- [17] K. B. Oldham and J. Spanier ; The Fractional Calculus, Academic Press (1974).
- [18] S. Moriguchi, K.Udagawa and S. Hitotsumatsu ; Mathematical Formulae, Vol.2, Iwanami Zensho, (1957), Iwanami, Japan.

Susana S. de Romero
CIMA, Facultad de Ingenieria
Universidad del Zulia
Apartado 10482
Maracaibo, VENEZUELA

Katsuyuki Nishimoto
Institute for Applied Mathematics
Descartes Press Co.
2 - 13 - 10, Kaguike, Koriyamy
963 - 8833 JAPAN